

Drug Information Sheet("Kusuri-no-Shiori")

Internal
Published: 12/2006

The information on this sheet is based on approvals granted by the Japanese regulatory authority. Approval details may vary by country. Medicines have adverse reactions (risks) as well as efficacies (benefits). It is important to minimize adverse reactions and maximize efficacy. To obtain a better therapeutic response, patients should understand their medication and cooperate with the treatment.

Brand name: Nyclin Powder 10%

Active ingredient: Nicotinic acid

Dosage form: white powder

Print on wrapping:



Effects of this medicine

This medicine is one of Vitamin B that work to maintain the metabolism of the body at a normal level, and acts to dilate blood vessels.

It is usually used for the prevention and treatment of various diseases that are caused by nicotinic acid deficiency including stomatitis, tinnitus, eczema, etc.

Before using this medicine, be sure to tell your doctor and pharmacist

- If you have previously experienced any allergic reactions (itch, rash, etc.) to any medicines.
- If you have hypotension.
- If you are pregnant or breastfeeding.
- If you are taking any other medicinal products. (Some medicines may interact to enhance or diminish medicinal effects. Beware of over-the-counter medicines and dietary supplements as well as other prescription medicines.)

Dosing schedule (How to take this medicine)

- Your dosing schedule prescribed by your doctor is << to be written by a healthcare professional >>
- In general, for adults, take 0.25 to 2 g (25 to 200 mg as the active ingredient) per day. The dosage may be adjusted according to the patient's age and symptoms. Strictly follow the instructions.
- If you miss a dose, skip the missed dose and continue your regular dosing schedule from the next dose onwards. You should never take two doses at one time.
- If you accidentally take more than your prescribed dose, consult with your doctor or pharmacist.
- Do not stop taking this medicine unless your doctor instructs you to do so.

Precautions while taking this medicine

Possible adverse reactions to this medicine

The most commonly reported adverse reactions include rash, lip swelling, cough, and shock-like symptom (facial pallor, cold sweat, dizziness on standing up). If any of these symptoms occur, consult with your doctor or pharmacist.

The symptoms described below are rarely seen as initial symptoms of the adverse reactions indicated in brackets. If any of these symptoms occur, stop taking this medicine and see your doctor immediately.

No pertinent entries.

The above symptoms do not describe all the adverse reactions to this medicine. Consult with your doctor or pharmacist if you notice any symptoms of concern other than those listed above.

Storage conditions and other information

- Keep out of the reach of infants and young children. Store away from direct sunlight, heat and humidity.
- Discard the remainder. Do not store them.

For healthcare professional use only / /

For further information, talk to your doctor or pharmacist.