Drug Information Sheet("Kusuri-no-Shiori")

Injection

Revised: 08/2023

The information on this sheet is based on approvals granted by the Japanese regulatory authority. Approval details may vary by country. Medicines have adverse reactions (risks) as well as efficacies (benefits). It is important to minimize adverse reactions and maximize efficacy. To obtain a better therapeutic response, patients should understand their medication and cooperate with the treatment.

Brand name:Cibenol Intravenous Injection 70mg

Active ingredient: Cibenzoline succinate

Dosage form: Injection

Imprint or print on wrapping:

Effects of this medicine

This medicine normalizes the pulse by blocking the sodium channel (membrane stabilizing activity) of cardiac muscle as well as prolonging the action potential duration.

It is usually used for the treatment of tachyarrhythmia.

The following patients may need to be careful when using this medicine.Be sure to tell your doctor and pharmacist.

・If you have previously experienced any allergic reactions (itch, rash, etc.) to any medicines or foods.

If you have atrioventricular block, sinoatrial block, congestive heart failure, angle-closure glaucoma or difficulty in urination.

If you are on dialysis.

・If you are pregnant or breastfeeding.

・If you are taking any other medicinal products. (Some medicines may interact to enhance or diminish medicinal effects. Beware of over-the-counter medicines and dietary supplements as well as other prescription medicines.)

Dosing schedule (How to take this medicine)

・Your dosing schedule prescribed by your doctor is((to be written by a healthcare professional))

・In general, this medicine is injected intravenously over 2 to 5 minutes. Blood pressure measurement and electrocardiography are performed on a continuous basis during treatment with this medicine.

・Consult with your doctor as to the specific treatment period.

Precautions while taking this medicine

Possible adverse reactions to this medicine

The most commonly reported adverse reactions include thirst, hot flush, palpitation, nausea, abdominal pain, numbness and chest pain. If any of these symptoms occur, consult with your doctor or pharmacist.

The symptoms described below are rarely seen as initial symptoms of the adverse reactions indicated in brackets. If any of these symptoms occur, see your doctor immediately.

・dizziness, chest pain, palpitation [proarrhythmic effect]

・pale face, cold sweat, dizziness on standing up [shock, cardiogenic shock, anaphylaxis]

・shortness of breath, general edema, cough [cardiac failure]

・general malaise, loss of appetite, yellowing of the skin and the whites of eyes [liver disorder due to circulatory failure, hepatic function disorder, jaundice]

The above symptoms do not describe all the adverse reactions to this medicine. Consult with your doctor or pharmacist if you notice any symptoms of concern other than those listed above.

Storage conditions and other information

For healthcare professional use only / /

For further information, talk to your doctor or pharmacist.