Drug Information Sheet("Kusuri-no-Shiori")

Internal
Revised: 04/2023

The information on this sheet is based on approvals granted by the Japanese regulatory authority. Approval details may vary by country. Medicines have adverse reactions (risks) as well as efficacies (benefits). It is important to minimize adverse reactions and maximize efficacy. To obtain a better therapeutic response, patients should understand their medication and cooperate with the treatment.

Brand name:Pronon Tablets 100mg

Active ingredient: Propafenone hydrochloride
Dosage form:white tablet (diameter: 7.6 mm, thickness: 3.5 mm)
Imprint or print on wrapping:(face) プロノン 100mg, 548, 不整脈治療剤
(back) プロノン 100mg, 不整脈治療剤

Effects of this medicine

This medicine works to normalize the pulse by blocking the sodium channel (membrane-stabilizing action) of cardiac muscle as well as prolonging the action potential duration.
It is usually used for the treatment of arrhythmia.

The following patients may need to be careful when using this medicine.Be sure to tell your doctor and pharmacist.

・If you have previously experienced any allergic reactions (itch, rash, etc.) to any medicines or foods.
If you have congestive heart failure, atrioventricular block and sinoatrial block.
・If you are pregnant or breastfeeding.
・If you are taking any other medicinal products. (Some medicines may interact to enhance or diminish medicinal effects. Beware of over-the-counter medicines and dietary supplements as well as other prescription medicines.)

Dosing schedule (How to take this medicine)

・Your dosing schedule prescribed by your doctor is((to be written by a healthcare professional))
・In general, for adults, take 150 mg as the active ingredient at a time, 3 times a day. The dosage may be adjusted according to the age or symptoms. Each tablet contains 100 mg of the active ingredient. Use this tablet when you take the medicine 100 mg or more at a time according to your dosing schedule described above. Strictly follow the instructions.
・If you miss a dose, take a dose as soon as possible. If it is almost time for the next dose, skip the missed dose and follow your regular dosing schedule. You should never take two doses at one time.
・If you accidentally take more than your prescribed dose, consult with your doctor or pharmacist.
・Do not stop taking this medicine unless your doctor instructs you to do so.

Precautions while taking this medicine

・Dizziness may occur after taking this medicine. Be aware of driving a car or operating machinery.
・Avoid taking any food containing hypericum perforatum (St. John's wort) since it may diminish medicinal effect.

Possible adverse reactions to this medicine

The most commonly reported adverse reactions include palpitation, nausea, vomiting, loss of appetite, abdominal pain, malaise, and muscle pain. If any of these symptoms occur, consult with your doctor or pharmacist.

The symptoms described below are rarely seen as initial symptoms of the adverse reactions indicated in brackets. If any of these symptoms occur, stop taking this medicine and see your doctor immediately.

・palpitation, shortness of breath, dizziness [ventricular tachycardia, ventricular fibrillation, sinoatrial block, atrioventricular block, syncope, sinus arrest, bradycardia]
・general malaise, loss of appetite, yellowness in the skin and/or whites of the eyes [liver dysfunction, jaundice]

The above symptoms do not describe all the adverse reactions to this medicine. Consult with your doctor or pharmacist if you notice any symptoms of concern other than those listed above.

Storage conditions and other information

・Keep out of the reach of infants and young children. Store away from direct sunlight, heat and humidity.
・Discard the remainder. Do not store them. If you do not know how to dispose of them, consult with your dispensing pharmacy or medical institution. Do not give this medicine to others.

For healthcare professional use only / /

For further information, talk to your doctor or pharmacist.