

Drug Information Sheet("Kusuri-no-Shiori")

Internal

Published: 06/2021

The information on this sheet is based on approvals granted by the Japanese regulatory authority. Approval details may vary by country. Medicines have adverse reactions (risks) as well as efficacies (benefits). It is important to minimize adverse reactions and maximize efficacy. To obtain a better therapeutic response, patients should understand their medication and cooperate with the treatment.

Brand name: Tadalafil Tablets 20mg AD "TE"

Active ingredient: Tadalafil

Dosage form: light red-brown tablet, diameter: 7.5 mm, thickness: 3.4 mm

Print on wrapping: (face) タダラフィル, AD, 20mg, 「TE」, 肺動脈性肺高血圧症のくすり
(back) タダラフィル 20mg AD「TE」, ニトログリセリンなどの硝酸薬と本剤は併用できません。



Effects of this medicine

This medicine relaxes blood vessels to lower pulmonary arterial pressure and pulmonary vascular resistance by inhibiting phosphodiesterase 5 (PDE5) enzyme in pulmonary vascular smooth muscle. It is usually used to treat pulmonary arterial hypertension (PAH).

Before using this medicine, be sure to tell your doctor and pharmacist

- If you have previously experienced any allergic reactions (itch, rash, etc.) to any medicines. If you have renal dysfunction, hepatic dysfunction, arrhythmia, hypotension, hypertension or retinitis pigmentosa. If you have a history of cerebral infarction/intracranial hemorrhage in these 6 months.
- If you are pregnant or breastfeeding.
- If you are taking any other medicinal products. (Some medicines may interact to enhance or diminish medicinal effects. Beware of over-the-counter medicines and dietary supplements as well as other prescription medicines.)

Dosing schedule (How to take this medicine)

- Your dosing schedule prescribed by your doctor is((to be written by a healthcare professional))
- In general, for adults, take 2 tablets (40 mg of the active ingredient) at a time, once a day. Strictly follow the instructions.
- If you miss a dose, take a dose as soon as possible. If it is almost time for the next dose, skip the missed dose and continue your regular dosing schedule. You should never take two doses at one time.
- If you accidentally take more than your prescribed dose, consult with your doctor or pharmacist.
- Do not stop taking this medicine unless your doctor instructs you to do so.

Precautions while taking this medicine

- The medicine may cause dizziness or vision disorder such as blurred vision. Be careful when working at heights, driving a car, or operating dangerous machinery.
- Avoid drinking grapefruit juice, as it may increase the effects and adverse reactions of the medicine.

Possible adverse reactions to this medicine

The most commonly reported adverse reactions include headache, flush, muscle pain and nausea. If any of these symptoms occur, consult with your doctor or pharmacist.

The symptoms described below are rarely seen as initial symptoms of the adverse reactions indicated in brackets. If any of these symptoms occur, stop taking this medicine and see your doctor immediately.

- rash, facial swelling, erosion in lips/mouth/conjunctiva [hypersensitivity]

The above symptoms do not describe all the adverse reactions to this medicine. Consult with your doctor or pharmacist if you notice any symptoms of concern other than those listed above.

Storage conditions and other information

- Keep out of the reach of children. Store away from direct sunlight, heat and moisture.
- Discard the remainder. Do not store them.

For healthcare professional use only / /

For further information, talk to your doctor or pharmacist.