

Drug Information Sheet("Kusuri-no-Shiori")

Injection

Published: 11/2019

The information on this sheet is based on approvals granted by the Japanese regulatory authority. Approval details may vary by country. Medicines have adverse reactions (risks) as well as efficacies (benefits). It is important to minimize adverse reactions and maximize efficacy. To obtain a better therapeutic response, patients should understand their medication and cooperate with the treatment.

Brand name: Trinosin-S injection 20 mg

Active ingredient: Adenosine triphosphate disodium hydrate

Dosage form: Injection

Print on wrapping:



Effects of this medicine

This medicine is involved in the physiological action in the living organisms as a high energy source. It also improves blood flow by dilating blood vessels.

It is commonly used for improvement of symptoms associated with late effects of head trauma, for the treatments of cardiac failure, muscular dystrophy and its related diseases, acute poliomyelitis, infantile cerebral palsy [flaccid], progressive spinal muscular atrophy and its related diseases, stabilization of accommodative function in accommodative asthenopia, tinnitus/deafness, or chronic gastritis accompanied by gastrointestinal hypomotility, or for improvement of liver function in chronic liver disease.

Before using this medicine, be sure to tell your doctor and pharmacist

- If you have previously experienced any allergic reactions (itch, rash, etc.) to any medicines.
If you are immediately after cerebral haemorrhage.
- If you are pregnant or breastfeeding.
- If you are taking any other medicinal products. (Some medicines may interact to enhance or diminish medicinal effects. Beware of over-the-counter medicines and dietary supplements as well as other prescription medicines.)

Dosing schedule (How to take this medicine)

- Your dosing schedule prescribed by your doctor is((to be written by a healthcare professional))
- In general, this medicine is used by intravenous injection, intravenous infusion, intramuscular injection or subcutaneous injection once or twice daily.

Precautions while taking this medicine

Possible adverse reactions to this medicine

The most commonly reported adverse reactions include feeling queasy/vomiting, loss of appetite, transient palpitations, and headache. If any of these symptoms occur, consult with your doctor or pharmacist.

The symptoms described below are rarely seen as initial symptoms of the adverse reactions indicated in brackets. If any of these symptoms occur, see your doctor immediately.

- Chest pain, feeling queasy, redness of face [shock-like symptom]

The above symptoms do not describe all the adverse reactions to this medicine. Consult with your doctor or pharmacist if you notice any symptoms of concern other than those listed above.

Storage conditions and other information

For healthcare professional use only / /

For further information, talk to your doctor or pharmacist.