

Drug Information Sheet("Kusuri-no-Shiori")

Internal
Revised: 04/2009

The information on this sheet is based on approvals granted by the Japanese regulatory authority. Approval details may vary by country. Medicines have adverse reactions (risks) as well as efficacies (benefits). It is important to minimize adverse reactions and maximize efficacy. To obtain a better therapeutic response, patients should understand their medication and cooperate with the treatment.

Brand name: Zeotin Tablets 100mg

Active ingredient: Methyl L-Cysteine hydrochloride

Dosage form: white tablet (φ : 9.5 mm, thickness: 5.7 mm)

Print on wrapping: (face) 301, ゼオチン 100mg, (back) ゼオチン, 100mg



Effects of this medicine

This medicine acts to reduce the viscosity of the sputum and pus and prompt the secretion of mucus from the respiratory tract, thereby easing the sputum discharge and nasal secretion.

It is usually used in patients with common cold, bronchial asthma, or acute and chronic bronchitis for expectoration, as well as patients with chronic sinusitis for pus discharge.

Before using this medicine, be sure to tell your doctor and pharmacist

- If you have previously experienced any allergic reactions (itch, rash, etc.) to any medicines.
- If you have liver disorder or cardiac failure.
- If you are pregnant or breastfeeding.
- If you are taking any other medicinal products. (Some medicines may interact to enhance or diminish medicinal effects. Beware of over-the-counter medicines and dietary supplements as well as other prescription medicines.)

Dosing schedule (How to take this medicine)

- Your dosing schedule prescribed by your doctor is << _____ to be written by a healthcare professional >>
- In general, for adults, take 1 tablet (100 mg as the active ingredient) at a time, 3 times a day. The dosage may be adjusted according to the patient's age and symptoms. Strictly follow the instructions. The tablet must not be chewed, but swallowed whole.
- If you miss a dose, skip the missed dose and continue your regular dosing schedule from the next dose onwards. You should never take two doses at one time.
- If you accidentally take more than your prescribed dose, consult with your doctor or pharmacist.
- Do not stop taking this medicine unless your doctor instructs you to do so.

Precautions while taking this medicine

Possible adverse reactions to this medicine

The most commonly reported adverse reactions include rash. If any of these symptoms occur, consult with your doctor or pharmacist.

The symptoms described below are rarely seen as initial symptoms of the adverse reactions indicated in brackets. If any of these symptoms occur, stop taking this medicine and see your doctor immediately.

No pertinent entries.

The above symptoms do not describe all the adverse reactions to this medicine. Consult with your doctor or pharmacist if you notice any symptoms of concern other than those listed above.

Storage conditions and other information

- Keep out of the reach of infants and young children. Store away from direct sunlight, heat and humidity.
- Discard the remainder. Do not store them.

For healthcare professional use only / /

For further information, talk to your doctor or pharmacist.